



Yours Energetically

by Jean-Pierre Petit


Translated by John Murphy























And in the daytime
I winch the counterweight
back up.

You're stocking POTENTIAL ENERGY.




It helps, but why is it always us that have to do the WORK?





I'm perfecting my method for STORING ENERGY.



The system that I've designed represents a storing of INTERNAL ENERGY





Sophie!
It was just the STORAGE
OF INTERNAL ENERGY!



CHEMICAL ENERGY



I'm going to tidy up the cave. Look at this saltpetre, sulphur...

And this charcoal left over from the forest fire that God Thunder made.







...just this big stone left







Sophie! I've found something.
There's ENERGY in this BLACK POWDER
I've just invented.




We'll be able to use it to cook things and keep warm!

You'll see...

Ok, but if you want my opinion, it isn't very easy to use.



Should I forget it?




That works!!!
The sand calms the mixture so that
it frees energy more slowly!




We won't be freezing cold this winter.





NUCLEAR ENERGY






...boxes with devils inside!?



An old legend said that once, ENERGY was shut inside the NUCLEII of certain ATOMS like URANIUM.

These atoms were made in the suns, in their infernal furnaces, then ejected and imprisoned in the mass of Earth when it was formed.

But these atoms aren't solid boxes and sometimes a top pops off.



The legend says that at the END OF TIME all the devils will be out of their boxes and there won't be any energy like this left in the universe.



But it will take a long time,




So how long wil the devils remain in their boxes? How long will the NUCLEII keep the ENERGY they have?



It depends on the boxes my son, it depends on the nucleii of the atoms.

RADIOACTIVE PERIOD OF AN ELEMENT

If we take an ensemble of boxes with devils in them, after a time T, called HALF-LIFE, or PERIOD, half of the devils will have been freed. In an identical lapse of time, half the remaing boxes will open in their turn, and so on. This half-life can vary greatly, from hundreds of thousands of years to a fraction of a second.



And if there weren't all these boxes with their devils, all these nucleii filled with energy inside the Earth then we'd be much colder in winter.

It would be nice if I could find all these atoms charged with energy.






If I could put enough of them into a bottle I could keep warm all winter.

Careful Archibald, the NUCLEAR ENERGY springs are much more powerful than CHEMICAL ENERGY, HUNDREDS OF THOUSANDS TIMES MORE POWERFUL.




So the devils emitted by the radioactive nucleii shoot out with great violence.







Good, after a time period equal to their half-life, half the boxes are empty.



After an equivalent lapse of time half the remaining boxes have opened and released their devils.





The Earth must have been a lot more radioactive in the beginning.


And then it calmed down.





CONVERSION OF ENERGY

But where's the HEAT in all that?



What if we put it in a cooking pot?





It works. The ENERGY emitted by the RADIOACTIVE ATOMS is absorbed by the water and CONVERTED INTO HEAT.



Yes but this NATURAL RADIOACTIVITY doesn't release much ENERGY.

So we'd need a lot of radioactive material to be able to keep ourselves warm.


THE DIFFERENT SPECIES OF DEVILS

Basically there is only one devil species. The first things that nucleii can emit are X or Y RADIATION. A sort of invisible light.





They can be absorbed with a sufficiently thick lead barrier and then their energy is converted into heat.


There are other types of devils that have an ELECTRIC CHARGE



Do they go fast?









NUCLEUS STABILITY




MESONS in the NUCLEII act a bit like ELECTRONS in MOLECULES: they ensure COHESION.



of molecules.





Electron ensuring
A MOLECULAR LIAISON.

CHEMISTRY interprets the rearrangement of MOLECULES.



NUCLEAR PHYSICS studies the REARRANGEMENT of NUCLEII A nucleus that is considered to be UNSTABLE is a nucleus with a short life


But neutrons, when acting on certain nucleii (themselves relatively stable having very long lives) can destabilise them completely and cause them to split, FISSION.







That's the case for URANIUM 235 and PLUTONIUM 239

FISSION





These nucleii can be represented as an assembly of two blocks of different masses and with one neutron.







Uranium 235 and Plutonium 239 nucleii have a certain type of natural radioactivity, associated with a very long period.










(*) This is a schematic image. In fact the incident neutron is first absorbed by the fuse nucleus (U235 becomes U236 and P239 becomes P240).


These are new objects, very unstable, which break down almost immediately.


CHAIN REACTIONS















CRITICAL CONDITIONS



In fact, between the weak NATURAL RADIOACTIVITY emission level and the CHAIN REACTION, we can find an average term. By adjusting this CONCENTRATION, which is quite difficult and delicate, we can set the number of devils that will be ejected each second, that is to say the energy flow.




NUCLEAR REACT




It looks like fly-paper.






The devils all are captured bit by bit.

There are practically no more chain reactions.




All that remains is "normal" energy emission, the natural energy of the radioactive body, which is considerably weaker.

So. To make a NUCLEAR REACTOR you just need to collect together enough heavy nucleii, URANIUM 235 or PLUTONIUM 239. And we can control the reactor's activity with a body that absorbs the devils, here they are FISSION neutrons.



And we'll use CADMIUM to absorb the NEUTRONS.


Apparently Plutonium 239 doesn't exist in nature so how can we think about using it in a reactor?




FERTILE MATERIAL

Uranium 238 could be considered to be an assembly of two elements. There's room left for a neutron.

In other words, when a
Uranium reactor is operating it
contains a mix of FISSILE material and FERTILE material. It
converts a certain amount of
the FERTILE material into
FISSILE material




it will change into Plutonium 239 and that is FISSILE



A certain amount? How much is that?

It depends how we operate the reactor. At first FISSION NEUTRONS are emitted in all directions, at 20,000 kilometers a second.




RAPID NEUTRON REACTORS

These RAPID NEUTRONS interact easily with fertile U238 thus creating Pu239 at a good rate.

What are you doing?



I'm loading my reactor with a mineral rich in Uranium 235 (enriched Uranium)




Then I put on a FERTILE COVER of U238



The FAST NEUTRONS move at 20,000 km/s in the HEART of the REACTOR. If we thought of them as gas molecules they would be at a temperature of 16 thousand million degrees.

THREE YEARS LATER

Oh! Archie has made MORE Pu239 than he has used U235, It's a SUPERGENERATOR




That's normal because each act of fission brings into play TWO rapid neutrons which allow the transformation of 2 U238 into Pu239


SLOW NEUTRON REACTORS

With CADMIUM I can absorb neutrons and so easily control the reactor's activity (or even stop it). But with GRAPHITE and HEAVY WATER I can SLOW the neutrons without absorbing them. These are called MODERATORS






In this way we can lower the THERMAL AGITATION SPEED of the neutorns to 2 Km/s. This neutron gas, cold, is at the general temperature of the reactor.



A little Pu 239 is still made but a lot less than in a rapid neutron reactor There is no clear frontier between the two types of reactor. There are also 'warm' reactors, half way between the two.



RADIOACTIVE WASTE INDUCED RADIOACTIVITY





The U235 and Pu239 nucleii break into two pieces in many different ways. Here the example is of Uranium 235 being split into radioactive Strontium 94 and Xenon 140. Note that 94+140+1=235

All that is a bit of a nuisance.

Many FISSION BY-PRODUCTS have a long life and remain radioactive for a very long time. STRONTIUM can fix itself in bone material and IODINE in the Thyroid.

Plutonium is also very dangerous, it can provoke CANCERS and LEUKEMIA





Fission neutrons can also be absorbed by peaceful atoms, such as those forming the structure of the reactor, which can make them dangerous and unstable as well as radioactive, so increasing the amount of radiactive waste.

MADE-TO-MEASURE RADIOELEMENTS



So a reactor produces unstable radioactive waste with different periods.


No, they're nucleii that are likely to lose mass by emitting Helium atoms, electrons or anti-electrons (*)

You mean that they're nucleii that are likely to split in their turn?



We can create 'made-to-measure' radioelements with different periods by placing certain elements in the reactor and submitting them to a bombardment of devils. In that way we get what is called artificial radioactivity.





A-BOMBS

Nuclear physics has allowed considerable progress to be made in fireworks science. In suddenly bringing together two masses of fissile material (U235 and Pu239) with the help of an explosive, we create critical conditions and provoke an intense chain reaction, with undeniable aesthetic effects.

Let's see. By bringing together these two masses I obtain a CRITICAL MASS

A great number of all sorts of devils are emitted and the radioactive waste rises into the upper atmosphere by ascendance which is caused by the release of intense heat. But that's nice because the neighbours can benefit too

It's coming, it's coming !..

If you want to join the HAPPY PYROTECHNICIANS you'll need a pure fissile material (100% U235 or Pu239). There are two ways to do it, either refine natural Uranium or apply to your neighbourhood reactor to collect the Pu239 that is produced after each cycle of operation.

FUSION


THE STATE OF THE S

Say, the Sun is a planet that must contain a lot of Uranium for it to be so hot.

No Archie, that isn't it.
In CHEMICAL REACTIONS we start of with a mix of substances, such as HYDROGEN and OXYGEN

But...nothing's happening ?!

It's because the temperature isn't high enough.














So there are lots of reactions that give off a great deal of energy without producing toxic substances.



If one day we use flying planes with a hydrogenoxygen mix (stored in liquid form) all they'd leave behind as they passed would be clouds!




Yes, if we brought their temperature up high enough



We could make DEUTERIUM react with TRITIUM, which are two sorts of HEAVY HYDROGEN (the hydrogen nucleus is light, constituted of a single proton P). The nucleii of these ISOTOPES are only differentiated by the number of neutrons they have. The mix od Deuterium and Tritium tends to give Helium

Here's an element of the gas HEAVY

GRAND DEVIL BALL



Deuterium molecule

Tritium molecule


31

Then the dance rhythm becomes really devilish. Molecules break up (disassociation) and the electron-bees orbit around a single nucleus.

APPROACHING THREE THOUSAND DEGREES


There's no way we can orbit around these nucelii, they're always moving.






Yep, it's becoming infernal. I give up...

The hot gas then becomes a soup of nucleii and free electrons, a HOT PLASMA.








In this case, FUSION is just as polluting as FISSION because these fusion neutrons will transform the nearby atoms, turning them into radioactive atoms.

So we try to absorb these neutrons with Lithium 6, which gives Helium 4 and Tritium 3.



HÉLIUM 4



TRITIUM 3


In other words, the envelope of Lithium 6 behaves as a "fertile" material. This reaction is supposed to give "fusion fuel", Tritium 3. Yes, a fusion reactor is related to a supergenerator. Luckily, because Tritium 3 is unstable (*) and doesn't exist in a natural state.





But only Tritium is regenerated.

Nevertheless, I see that there are all sorts of fusion reactions and nucleii rearrangements which don't release free neutrons.






Lithium 7 + Hydrogen 1 (light) gives
2 Helium 4

$$(7 + 1 = 2 \times 4)$$


Boron 11 + Hydrogen 1 gives 3 Helium 4

$$(11 + 1 = 3 \times 4)$$




The first has an ignition temperature of 500 million degrees and, for the second, close on a thousand million degrees!..

Hmm...obviously...but exactly how do we fuse these nucleii?




In the heart of the sun it happens slowly but at a temperature that is only 15 million degrees.





Yes, to get a nuclear "fire" you need 150 million degrees for reactions to take place in, let's say, a period of time of the order of a second.





(*) Edward teller, researcher at Los Alamos during the war, was the model for Doctor Strangelove in the film 'How I stopped worrying and learned to love the bomb'.





But of course, think about it.

The H-bomb exploded. Neutrons from fusion attacked the FERTILE U238 material and transfromed it into Pu239 which fissioned immediately

That was the terrible FISSION-FUSION-FISSION bomb

FUSION BY DIRECTED ENERGY

An attempt was made to create FUSION by localising onto a DEUTÉRIUM-TRITIUM mixture (in liquid state) all the energy forms: radiations, emitted by powerful LASERS, various particles: electrons, nucleii from accelerators. The POWER needed is phenomenal. To set off this THERMONUCLEAR fire, an energy equivalent to that of a solar mirror the size of France needs to be concentrated (for a few thousand millionths of a second) on a sphere of 1 mm diameter

Yep, I'm

getting a tan.

The INSTANT POWER is enormous but the global ENERGY remains modest: this nuclear "match" is equivalent to two hundred grams of powder.

EPILOGUE

We need NUCLEAR ENERGY. But all that, FUSION, FISSION has many disadvantages

The awkward waste for instance.



And plenty of risks
of accidents. If a reactor starts to
overheat it will melt the steel and
concrete container, even the floor
(CHINESE SYNDROME (*)) and the
mass in fission will force itself
down into the ground without our
being able to stop it.

40 years isn't very long. We are only at the beginning of the NUCLEAR ERA.




I believe in the possible revolutionary progress that can be made, which could completely change the basic problem, but more on the FUSION side than that of FISSION.



(*) An image due to atomists, according to whom the reactor going through the Earth from one side to the other would reappear in... China!...

Theoretically, in fusion reactions, where free neutrons don't intervene, we could CONFINE the FUSION PLASMAS by using powerful magnetic equipment (charged particlesd "run away" from regions with intense magnetic fields).


THE GOLDEN AGE! The fusion generating station, non-polluting (lithium-hydrogen or boron-hydrogen). The only product of the reaction is helium which we could use to inflate balloons.



Don't make me laugh, it's a dream!

Nevertheless, catalytic stoves exist that allow heat to be produced AT HOME with the windows closed and without using a chimney.

It's true, it creates water vapour and carbon dioxide which can be breathed in in moderate quantities.



Could a FUSION CATALYSER exist that would allow operation at a suitably low temperature?



We already know one: Carbon

Ah yes, in fact, how does the Sun manage to work by fusion when its central boiler is at 15 MILLION DEGREES, that is to say at a temperature TEN TIMES LESS THAN THE IGNITION TEMPERATURE which is 150 MILLION DEGREES?

The the stain the

The Carbon serves as a catalyst. It intervenes in the stages, fairly complex ones, of the reaction and, in the end, is regenerated. It begins with Carbon 12 then Hydrogen 1 giving Nitrogen 13.

Then the Nitrogen 13 is transformed into Nitrogen 15 and finally: Nitrogen 15 + Hydrogen 1 -> Carbon 12 + Helium 4 (Bethe's cycle).

But this reaction is far too SLOW (except for the Sun, which has plenty of time).

MUONS

We can create complex chemical reactions in a cold gaseous mix by bombarding the molecules with electrons via a simple electrical discharge.


For example:

2CH4 (methane) + 2

gives C2H2 (acetylene)

+3H2

In a molecule we can replace the electrons with MUONS, particles which resemble big electrons and which bring the various nucleii closer together.







NO PROBLEM SIR. We know how to create muons in an accelerator. When they hit deuterium and tritium nucleii it creates helium. Therefore there is fusion. But between this experiment in microphysics, which just concerns a few particles, and a usable industrial fusion there is quite a distance to travel!!!

We could also play with the SPINS of nucleii. That is to say make them dance a waltz instead of a tango. It improves the efficacity of the collisions.









All this is just beginning.
What do you think Archibald?

